



Contents lists available at ScienceDirect

Journal of Cranio-Maxillo-Facial Surgery

journal homepage: www.jcmfs.com

Surgical management of upgaze diplopia in patients after posttraumatic orbital floor reconstruction

Piotr Loba ^{a, *}, Marcin Kozakiewicz ^b, Anna Broniarczyk – Loba ^c^a Department of Ophthalmology (Head: Wojciech Omulecki), Medical University of Lodz, Kopcinskiego 22, Lodz 90-153, Poland^b Department of Maxillofacial Surgery (Head: Marcin Kozakiewicz), Medical University of Lodz, Zeromskiego 113, Lodz 90-549, Poland^c Department of Binocular Vision Pathophysiology and Strabismus (Head: Anna Broniarczyk-Loba), Medical University of Lodz, Kopcinskiego 22, Lodz 90-153, Poland

ARTICLE INFO

Article history:

Paper received 7 January 2015

Accepted 27 March 2015

Available online xxx

Keywords:

Orbital fracture

Diplopia

Reconstruction surgery

Strabismus

ABSTRACT

Objective: The most common complication of otherwise successful reconstructive surgery of a fractured orbital floor is persistent diplopia. For patients with troublesome double vision in upgaze, a reasonable solution is offered by strabismus surgery. The aim of our study is to examine the results of extraocular muscle surgery in cases of diplopia that persisted in upgaze after posttraumatic orbital floor reconstruction. **Material and methods:** In this study we present a retrospective series of 24 patients with troublesome vertical diplopia in upgaze. In all cases, the surgery consisted of a posterior fixation suture placement on the contralateral superior rectus muscle with or without its recession. Full orthoptic examination was conducted before and 3 months after the surgery.

Results: Postoperatively 19 patients (79%) were diplopia free and 6 (21%) had vertical diplopia in extreme upgaze. The field of binocular single vision improved threefold. None of the patients reported diplopia in the primary position or in any position other than upgaze.

Conclusion: Vertical inconstant strabismus and diplopia in upgaze persisting after orbital reconstructive surgery may be corrected surgically. Contralateral posterior fixation of the superior rectus muscle, with or without its recession, appears to be an effective procedure for use in these patients.

© 2015 European Association for Cranio-Maxillo-Facial Surgery. Published by Elsevier Ltd. All rights reserved.

1. Introduction

The most common complication of otherwise successful reconstructive surgery of a fractured orbital floor is persistent diplopia (Biesman et al., 1996), a condition that affects 5%–18% of patients who have undergone this surgery (Kerr, 2004; Ceylan et al., 2011; Loba et al., 2012). Usually double vision is experienced during upgaze, extending over various areas and sometimes into primary position. Diplopia is less common in other gaze directions (Morax and Pascal, 1984; Gosse et al., 2010).

Although some cases require a revision of the orbital floor by a maxillofacial surgeon (Kozakiewicz and Szymor, 2013), no such procedure is indicated in others in whom no implant malposition or persistent herniation of orbital tissue into the maxillary sinus is revealed by postoperative imaging. The cause of ocular

misalignment in such patients is known to be ipsilateral hypotropia, which in turn is caused by a restricted inferior rectus (Biesman et al., 1996). This incomitance is greatest in upgaze but is sometimes also present in primary position. The typical treatment for such patients is a period of observation for spontaneous improvement. However, for those patients who experience sustained diplopia in upgaze, a reasonable solution is offered by strabismus surgery, the goal of which is to broaden the field of binocular single vision and to improve ocular alignment in upgaze without affecting other gaze directions.

The aim of our study is to examine the results of such extraocular muscle surgery in cases of diplopia that persisted in upgaze after posttraumatic orbital floor reconstruction.

2. Materials and methods

During a 6-year period, 264 patients were referred to the Strabismus Outpatient Clinic with residual diplopia and abnormal ocular motility following reconstructive surgery for orbital fracture.

* Corresponding author. Tel.: +48 606 990050; fax: +48 42 6776800.

E-mail addresses: ploba@onet.pl (P. Loba), marcin.kozakiewicz@umed.lodz.pl (M. Kozakiewicz), abl1@op.pl (A. Broniarczyk – Loba).

<http://dx.doi.org/10.1016/j.jcms.2015.03.034>

1010-5182/© 2015 European Association for Cranio-Maxillo-Facial Surgery. Published by Elsevier Ltd. All rights reserved.

Routinely a 6-month observation period was carried out for spontaneous resolution of symptoms. Patients suspected of failed tissue release or implant malposition underwent a postoperative computed tomographic (CT) scan and were referred back to the maxillo-facial surgeon for reoperation. Such patients were further observed postoperatively.

After the observational period, 33% (87) of the patients still experienced diplopia in various gaze directions. In most cases, diplopia was limited only to the edges of the peripheral gaze and did not affect daily activities. Troublesome double vision in upgaze was reported by 30 patients to whom surgical treatment was proposed. Of these patients, 24 consented to the surgery (17 male and 7 female; mean age, 39.6 ± 12.4 years). A more detailed profile of the characteristics of the group is given in Table 1.

Orthoptic assessment was performed before surgery and 3 months postoperatively in each patient. Ocular rotations, the type and direction of diplopia, and the vertical angle of strabismus measured by prism and cover test were noted in prism diopters [Δ]. Hess charts were plotted in each case as well as Binocular Single Vision Field (BSV) (Medmont M900). A passive duction test was performed in the theater under general anesthesia.

The type of surgery was single-step and similar in all cases. In order to restore vertical comitance in upgaze, the procedure of choice was placement of a 14-mm posterior fixation suture on the contralateral superior rectus muscle. In 14 cases (58%), a small recession was added. The amount of recession was adjusted in accordance with the angle of deviation in primary position (none up to 3Δ ; 2 mm, 3Δ to 5Δ ; 3 mm, 6Δ to 14Δ ; 4 mm, 15Δ and more).

The results were subject to statistical analysis with Statistica 10.0, using a t-test for paired measurements ($p < 0.01$). The methods applied in the study adhered to the Declaration of Helsinki and were accepted by the Board of Ethics of the Medical University of Lodz (RNN/144/09/KE).

3. Results

During surgery, a restriction of inferior rectus muscle on the side of the injury was confirmed by passive force duction test in all cases. No major complications occurred during the procedure in any patient. No upper lid retraction was noted, but prolonged lid swelling (up to 5 days) was reported after the surgery in 4 cases.

Before the surgery, the mean vertical deviation in upgaze was found to be $19.6 \pm 8.9\Delta$, ranging from 10Δ to 45Δ , in primary position $4.99 \pm 0.78\Delta$, ranging from 0Δ to 16Δ , and $1.20 \pm 2.17\Delta$ in downgaze, ranging from -1.5Δ to 7Δ . When comparing the orthoptic measurements obtained before and 3 months after surgery, a significant reduction of mean VD was revealed in upgaze ($p = 0.000$). Vertical deviation in the primary position changed significantly ($p = 0.000$) only in patients who underwent additional

recession of the superior rectus. Mean recession was 2.43 ± 0.62 mm correcting $2.9 \pm 0.5\Delta/\text{mm}$. Mean angles of deviation and BSV before and after surgery are summarized in Table 2.

The rate of success, understood as a vertical angle of less than 5Δ in upgaze, was 66.5%. Undercorrections were noted in six cases (21%) and overcorrections in three (12.5%). Subjective evaluation showed that 19 patients (79%) were diplopia free and 6 (21%) had vertical diplopia in extreme upgaze. The field of binocular single vision (BSV) improved threefold with a mean postoperative BSV loss of less than 10%. Postoperatively, none of the patients reported diplopia in the primary position or in any position other than upgaze. An example of the ocular alignments and Hess charts before and after surgery is presented in Figs. 1 and 2.

4. Discussion

Orbital reconstruction surgery is intended to correct cases of limited elevation of the affected eye, a hallmark of orbital floor fractures, by freeing muscles and connective tissue from the fracture and repositioning them from the maxillary sinus. Nevertheless, although reconstructive surgery may appear to be carried out correctly, vertical incomitance and thus diplopia may still persist in some cases. The cause of diplopia may be attributed to the presence of tissue adhesions, implant malposition, or improper tissue reposition (Kushner, 1982; Van Eeckhoutte et al., 1992; Biesman et al., 1996). Computed tomographic imaging is sometimes helpful in identifying the problem (Jazwiecka-Koscielniak and Kozakiewicz, 2014). However there is often no apparent cause, and the patient is left to await the spontaneous resolution of symptoms. A 6-month waiting period is generally accepted as sufficient time for such spontaneous recoveries to occur, especially in small deviations (Van Eeckhoutte et al., 1992). In the present study, 27% of patients with persistent diplopia required eye muscle surgery, a proportion similar to that observed previously (Loba et al., 2012).

Generally, there are several approaches to improving ocular alignment and diplopia in patients with incomitant vertical strabismus. Ipsilateral inferior rectus recession, followed if necessary by resection of the superior rectus on the same side, is recommended by Van Eeckhoutte et al. (1998), who report good results in all but one patient, who was overcorrected. Kerr also describes recession of the inferior rectus, but with the use of adjustable sutures, resulting in an alignment in primary position and improvement in upgaze. However, the study notes that large recessions of the inferior rectus might negatively influence depression and thus should be linked with contralateral inferior rectus recession. An alternative procedure advocated by the author is based on a recess–resect procedure on the affected side to split the amount of surgery to both vertical recti (Kerr, 2004).

After reconstructive surgery for orbital floor fractures, the inferior rectus muscle on the side of the injury is usually fibrously changed and its function is impaired. Some weakness of the inferior

Table 1
Selected group characteristics (n = 24).

Type of fracture	
Orbital floor	24 (100%)
Medial wall	7 (29%)
Orbital rim involvement	11 (46%)
Number of orbital surgeries	
One	20 (83%)
Two	4 (17%)
Type of implant	
Polypropylene sheet	17 (71%)
Titanium mesh	5 (21%)
Bone graft	2 (8%)
Time from trauma to first reconstruction	
14th day	10 (42%)
>14 days	14 (58%)

Table 2
Mean angles of deviation before and after the surgery.

Angle in diopters (Δ)	Before surgery	After surgery	p-Value
Upgaze	19.6 ± 8.9	2.58 ± 2.97	0.000
Primary position (recession group)	8.38 ± 3.96	1.18 ± 1.81	0.000
Primary position (no recession group)	0.25 ± 0.26	0.22 ± 0.25	0.797
Downgaze	1.20 ± 2.17	0.52 ± 0.95	0.129
BSV loss [%]	40.3 ± 16.5	9.4 ± 6.3	0.000

BSV, loss of binocular single vision field (%).




Fig. 1. Ocular alignment in vertical gaze and Hess chart before surgery on the extraocular muscles in an exemplary patient. Trauma to the right orbit.

rectus can be noted in every case of its restriction, which can be demonstrated by careful assessment of ocular alignment in downgaze, during which a small amount of ipsilateral hypertropia may be noted. Similarly, patients presenting with diplopia in upgaze also demonstrate a common pattern of simultaneous elevation and depression deficit (Morax and Pascal, 1984; Gosse et al., 2010; Loba et al., 2012). The exact etiology of this phenomenon is not well established but it might be considered as a specific variant of inferior rectus restriction (Lerman, 1970).

Therefore, the amount of recession of ipsilateral inferior rectus muscle that should improve elevation of the eye but not lead to postoperative diplopia in downgaze is difficult to establish preoperatively. Recession of this already weaker muscle can result in its underaction, thus requiring subsequent, additional surgery (Kushner, 1982). Some procedures can be used to restore comitance

in downgaze, such as posterior fixation of the contralateral inferior rectus, suggested by Buckley and Meekins, or plain recession, as suggested by Kerr (Buckley and Meekins, 1988; Kerr, 2004).

Our approach eliminates the above problems. The surgery does not affect downgaze and is a one-step procedure. The idea of using posterior fixation sutures in vertical incomitancies is not new. Kouri et al. reported 34 cases, among which 5 were blow-out fractures (Kouri et al., 2002). A similar surgical procedure was used by Hoover, although in single vertical rectus palsies rather than blow-out fracture cases, and good results were obtained with the use of adjustable sutures for recession (Hoover, 1998).

The influence on primary position is regulated by additional recession and should be aimed at around 3Δ improvement per millimeter. However, the surgeon should be aware of possible overcorrection leading to reversal in incomitance: the operated eye




Fig. 2. Ocular alignment in vertical gaze and Hess chart after posterior fixation (14-mm) with 3-mm recession of the left superior rectus muscle. Trauma to the right orbit.

becomes hypotropic. In our opinion, it is better to slightly undercorrect in such cases, as diplopia remaining in upgaze is tolerable. On the other hand, all our undercorrections occurred in patients with preoperative large angles and diplopia extending over the primary position as far as downgaze.

5. Conclusion

In conclusion, contralateral posterior fixation of the superior rectus muscle, with or without its recession, appears to be an effective and stable procedure for patients with vertical incomitant deviation and diplopia in upgaze persisting after orbital reconstruction surgery. The described method does not cause the postoperative hypertropia or diplopia in downgaze that might appear after ipsilateral recession of the inferior rectus muscle.

Financial support

No financial support.

Conflict of interest statement

The authors certify that there is no conflict of interest with any financial organization regarding the material discussed in the manuscript.

Acknowledgments

The authors would like to thank the team from the Department of Cranio-Maxillo-facial and Oncological Surgery from Medical University of Lodz for their support in gathering the material for this study.

References

- Biesman BS, Hornblass A, Lisman R, Kazlas M: Diplopia after surgical repair of orbital floor fractures. *Ophthalmol Plast Reconstr Surg* 12: 9–16, 1996
- Buckley EG, Meekins BB: Faden operation for the management of complicated incomitant vertical strabismus. *Am J Ophthalmol* 105: 304–312, 1988
- Ceylan OM, Uysal Y, Mutlu FM, Tuncer K, Altinsoy HI: Management of diplopia in patients with blowout fractures. *Indian J Ophthalmol* 59: 461–464, 2011
- Gosse EM, Ferguson AW, Lymburn EG, Gilmour C, MacEwen CJ: Blow-out fractures: patterns of ocular motility and effect of surgical repair. *Br J Oral Maxillofac Surg* 48: 40–43, 2010
- Hoover DL: Results of a combined adjustable recession and posterior fixation suture of the same vertical rectus muscle for incomitant vertical strabismus. *J AAPOS* 2: 336–339, 1998
- Jazwiecka-Koscielniak E, Kozakiewicz M: A new modification of the individually designed polymer implant visible in X-ray for orbital reconstruction. *J Craniomaxillofac Surg* 42: 1520–1529, 2014
- Kerr NC: Management of strabismus following orbital fracture repair. *Am Orthot J* 54: 24–31, 2004
- Kouri AS, Bessant DA, Adams GG, Sloper JJ, Lee JP: Quantitative changes in the field of binocular single vision following a fadenoperation to a vertical rectus muscle. *J AAPOS* 6: 294–299, 2002
- Kozakiewicz M, Szymor P: Comparison of pre-bent titanium mesh versus polyethylene implants in patient specific orbital reconstructions. *Head Face Med* 9: 32, 2013
- Kushner BJ: Paresis and restriction of the inferior rectus muscle after orbital floor fracture. *Am J Ophthalmol* 94: 81–86, 1982
- Lerman S: Blowout fracture of the orbit. Diagnosis and treatment. *Br J Ophthalmol* 54: 90–98, 1970
- Loba P, Kozakiewicz M, Nowakowska O, Omulecki W, Broniarczyk-Loba A: Management of persistent diplopia after surgical repair of orbital fractures. *J AAPOS* 16: 548–553, 2012
- Morax S, Pascal D: Surgical treatment of diplopia caused by fractures of the orbital floor. *J Fr Ophtalmol* 7: 633–647, 1984
- Van Eeckhoutte L, Apers R, De Clippeleir L, Van Lammeren M, Janssens H: Surgery on the extra-ocular muscles after orbital fracture. *Bull Soc Belge Ophtalmol* 245: 53–59, 1992
- Van Eeckhoutte L, De Clippeleir L, Apers R, Lammeren M, Janssens H, Baekeland L: A protocol for extraocular muscle surgery after orbital floor fracture (“blow-out”). *Binocul Vis Strabismus Q* 13: 29–36, 1998